

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.

Lorsban®-4E

EPA Reg. No. 62719-220

EPA 24(c) Special Local Need Registration SLN ID-010004
For Distribution and Use Only in the State of Idaho

RECEIVED

AUG 19 2005

DIV. OF AG. RESOURCES

Control of Grape Mealybugs and Cutworms Infesting Grapes

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Lorsban®-4E insecticide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of Lorsban-4E according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for Lorsban-4E.

Directions for Use

Cutworms: Apply Lorsban-4E at the rate of 1 quart per acre as a broadcast spray in a minimum spray volume of at least 50 gallons of water using power-operated ground spray equipment. Treat when cutworms first become active and when field counts indicate damaging insect populations are developing or present. Do not apply after bloom stage of growth. Consult your state agricultural experiment station or extension service specialist concerning cutworm control practices in your area.

Grape Mealybugs: Apply Lorsban-4E at the rate of 1 to 2 quarts per acre in a minimum spray volume of at least 50 gallons of water per acre using ground equipment only. Use the 1 quart rate for normal infestations prior to bloom stage of growth. Use the 2 quart rate for high infestations as a dormant or late dormant application only.

Specific Use Restrictions:

- **Preharvest Interval:** Do not apply within 90 days of harvest.
- Do not apply more than 2 quarts per acre of Lorsban-4E per crop growing season or make more than 2 applications.
- **Chemigation:** Do not apply through any type of irrigation equipment.
- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Specific Use Precautions:

- Applications after budbreak may result in transient leaf yellowing (Concords).

®Trademark of Dow AgroSciences LLC

R062-012

Approved: __/__/__

Replaces D06-062-030

Restrictions: Added "Specific Use" to "Restrictions" and "Precautions"; deleted heading "Use and Dosage Recommendations"; revised label with IRED-mandated language.